

## **TagFP Selector**

### **Data Sheet**

#### TagFP Selector

Catalog No.	N0510
Resin µL	1000
Slurry µL	2000

#### Description

TagFP Selector is based on a high-affinity single-domain antibody (sdAb) that is covalently immobilized on 4 % cross-linked agarose beads. The innovative, oriented and selective attachment via a flexible linker guarantees a high accessibility of the sdAbs and largely eliminates batch-to-batch variations. Due to the single-chain nature of sdAbs and their stable and covalent attachment, no leakage of light and heavy chains is observed during elution with SDS sample buffer. TagFP Selector thus features high affinity and superior capacity for TagFP fusion proteins while showing negligible unspecific background.

TagFP Selector is compatible not only with physiological buffers but also with high stringency buffers (see "Buffer Compatibility" below). TagFP Selector thus provides great freedom to adjust the binding and washing conditions to the experimental needs.

For recommended protocols please see our webpage at www.nano-tag.com/protocols.

To be used in vitro / for research only, not for diagnostic or therapeutic use! Non-toxic, non-hazardous, non-infectious.



## **TagFP Selector**

Properties				
Support	4 % cross-linked agarose, bead size 50-150 $\mu m$			
Coating	sdAb anti-TagFP clone 1H7			
Reactivity	Recognizes mTagBFP, mTagRFP, mKate and most common fluorescent proteins deriving from <i>Entacmaea quadricolor.</i>			
	Does not cross-react with common GFP- or dsRed/RFP derivatives.			
	For a detailed overview visit our webpage at www.nano-tag.com/protocols.			
Capacity	> 4 µg mTagBFP per µl of packed beads			
Formulation	50 % slurry in PBS containing 20 % Ethanol			
Shipment	Shipped at ambient temperature			
Storage	Store at 4 °C, do not freeze			
Stability	Stable for 12 months			

Buffer

- · Common buffer substances at pH 5 to 9
- Compatibility 2% Triton X-100, 1% Tween-20, 1% NP-40, 1% CHAPS, 1% Deoxycholate, 0.1% SDS
  - 4 M NaCl, 2 M KCl, 1 M MgCl<sub>2</sub>, 100 mM EDTA
    - 4 M urea
    - 10 mM DTT, 10 mM 2-Mercaptoethanol
    - RNAse A, DNAse I, Benzonase, protease inhibitors

For more information please visit our web page at www.nano-tag.com

#### Limited Use Label License

The purchaser of this product is granted a limited, non-transferable right to use the purchased material only for internal, academic research purposes. The license explicitly excludes the right to resell this product or any of its components. For information on further licensing possibilities please contact info@nano-tag.com

NanoTag Biotechnologies GmbH, Rudolf-Wissell-Str. 28a, 37079 Göttingen, Germany Phone: +49 551 50556-365, E-mail: info@nano-tag.com, Web: www.nano-tag.com



# **TagFP Selector**

Fluorescent Protein	<b>∢</b> GFP Selector	RFP Selector	TagFP Selector
Sirius			
tSapphire	~		
Cerulean			
eCFP	~		
mTurquoise, mTurquoise2	7		
acGFP	~		
EGFP	,		
Emerald GFP	~		
mEGFP (A206K)	7		
mEGFP (L221K)	~		
superecliptic pHluorin			
paGFP	~		
superfolder GFP	<i>V</i>		
eYFP	<i>y</i>		
mVenus	7		
Citrine	•		
mRFP		<i>'</i>	
mCherry		<b>V</b>	
mScarlet		<b>V</b>	
mOrange2		<b>V</b>	
tdTomato		~	
dsRed1 , dsRed2		~	
mTagBFP			•
mKate, mKate2			<b>V</b>
mTagRFP			<b>V</b>
mTagRFP657			•

Not recognized: Dendra2, Dronpa, tdEOS, mEOS3.2, mRuby3 and mTFP (mTeal)